

# WHEN SHOULD I SEND MY CHILD TO SCHOOL?

Symptoms and Illnesses	Should My Child Go To School?
<b>Parent is Sick, Stressed, Hospitalized</b>	<b>YES</b> - If you are sick, your child still needs to attend school. Your illness does not excuse your child from attending. We all are sick at times so plan ahead for these days. Get a neighbor, relative or spouse to take your child to school and pick him or her up.
<b>Chronic Diseases/Illnesses (Asthma, Diabetes, Sickle Cell, Epilepsy etc.)</b> Chronic disease is a long-lasting condition that can be controlled but not cured.	<b>YES</b> – Your child should attend school. School personnel are trained to assist your child with his or her chronic disease and associated needs.
<b>Child Doesn't Want to go to School</b> Frequent crying, fear, anger, not wanting to socialize, behavior change, stomach ache, nausea (These can be signs of depression, anxiety, post-traumatic stress, or fear)	<b>YES</b> – You should keep your child in school, but try to determine what is causing the changes. Talk to school personnel and consult a health care provider. Your child may be experiencing bullying or trauma, may be behind in his or her school work or not getting along with others. These and other issues may require your or school personnel's attention.
<b>Cold Symptoms</b> Stuffy nose/runny nose, sneezing, mild cough	<b>YES</b> - If your child is able to participate in school activities send him or her to school.
<b>Conjunctivitis (Pink Eye)</b> The white of the eye is pink and there is a thick yellow/green discharge.	<b>YES</b> – Contact your health care provider to determine if treatment is necessary. Your child can attend school, when they are no longer contagious (24 hours after beginning treatment)).
<b>Head Lice</b> Intense itching of the head; may feel like something is moving	<b>YES</b> – Once you have been notified that your child has lice you must complete an initial treatment of shampooing of hair with a product for lice. Your child should be back to school the next day.
<b>Strains, Sprains and Pains</b>	<b>YES</b> – If there is no known injury and your child is able to function (walk, talk, eat) he or she should be in school. If pain is severe or doesn't stop, consult a health care provider.
<b>Menstrual Issues</b>	<b>YES</b> – Most of the time menstrual (periods) issues should not be a problem. If they are severe and interfering with your daughter attending school, consult with a health care provider.
<b>Fever</b> Fever usually means illness, especially if your child has a fever of 101 or higher as well as other symptoms like behavior change, rash, sore throat, vomiting etc.	<b>NO</b> – If your child has a fever of 100 or higher, keep them at home until his or her fever is below 100 for 24 hours without the use of fever reducing medication. If the fever does not go away after 2-3 days or is 102.0 or higher, you should consult a health care provider.
<b>Diarrhea</b> Frequent, loose or watery stool may mean illness but can also be caused by food and medication	<b>NO</b> – If, in addition to diarrhea, your child acts ill, has a fever or is vomiting, keep him or her at home. If stool is bloody, if the child has abdominal pain, fever or vomiting, you should consult a health care provider.
<b>Vomiting</b> Child has vomited 2 or more times in a 24 hour period	<b>NO</b> – Keep your child at home until the vomiting has stopped for 24 hours. If vomiting continues, contact a health care provider.
<b>Coughing</b> Severe, uncontrolled, rapid coughing, wheezing, or difficulty breathing	<b>NO</b> – Keep your child home and contact a health care provider. Asthma - if symptoms are due to asthma, provide treatment according to instructions given to you by your child's health care provider. When symptoms are controlled send your child to school.
<b>Rash with and without Fever</b>	<b>NO</b> – You must provide a note from a health care provider with a diagnosis of the cause of the rash. We will follow guidelines provided by the health department to determine when your child can return to school.
<b>Strep Throat</b> Sore throat, fever, stomach ache, and red, swollen tonsils	<b>NO</b> – Keep your child at home for the first 24 hours after an antibiotic is begun.
<b>Vaccine Preventable Diseases</b> <b>Chicken Pox</b> - fever, headache, stomach ache or sore throat, then a red itchy skin rash develops on the stomach first and then limbs and face. <b>Measles &amp; Rubella (German Measles)</b> – swollen glands, rash that starts behind ears then the face and the rest of the body, sore joints, mild fever and cough, red eyes <b>Mumps</b> – fever, headache, muscle aches, loss of appetite, swollen tender salivary glands <b>Pertussis (Whooping Cough)</b> – many rapid coughs followed by a high-pitched “whoop”, vomiting, very tired	<b>NO</b> – Keep your child at home until a health care provider has determined that your child is not contagious. We will follow the guidelines for communicable diseases from the local health department.  For questions, call your school nurse.

## DUNSMUIR JOINT UNION HIGH SCHOOL DISTRICT

5805 High School Way, Dunsmuir, California 96025 [www.dunsmuirhigh.k12.ca.us](http://www.dunsmuirhigh.k12.ca.us)

Source: <http://atschool.alcoda.org/files/healthflyer2013.pdf>, February 6, 2017

03/2017, pm, SCOE